

DNA Damage Processing at Telomeres: Maintaining Youthful Chromosomes




Patty O'presko




Patricia Opresko Dept EOH, Graduate School of Public Health
University of Pittsburgh Cancer Institute, Hillman Cancer Center

Relationship Between Telomere Length, Aging and Human Disease



(TTAGGG)₁₅₀₀



Diseases

- osteoporosis
- cancer
- heart disease/ atherosclerosis
- arthritis/ ulcerative colitis
- pulmonary fibrosis/ liver cirrhosis
- diabetes
- neurodegeneration

Factors

- perceived stress
- smoking/ polycyclic aromatic HC
- obesity
- environmental metals
- lack of exercise
- diet


Genomic instability and the loss of cell viability induced by telomere dysfunction contribute to disease and tissue/organ decline with age

Mechanisms of Telomere Loss and Preservation

Significance

Understanding mechanisms of telomere loss and preservation could inform about new therapeutic strategies that:

- 1) Protect telomeres in healthy cells to prevent telomere dysfunction-induced pathologies
 - cell senescence and degenerative disease associated with aging
 - genomic instability and carcinogenesis
- 2) Deplete telomeres to halt cancer cell proliferation
 - combine with telomerase inhibitors




Outline

General background on telomere biology
DNA damage and repair

1. Impact of bulky DNA lesions on telomere replication
2. Repair of bulky DNA lesions at telomeres

Summary and Conclusions

Roles for Dysfunctional Telomeres in Cancer and Aging




Somatic cells:
most have no telomerase activity


Stem cells:
variable amounts

Cancer cells:
90% express high telomerase activity


Telomere Cap is a Complex of Proteins and DNA Structure



Griffith et al
1999, Cell



Telomere Uncapping



Conditional knock out of TRF2 in mouse embryonic fibroblasts.

p53 proficient – senescence or apoptosis

p53 deficient – chromosome fusions

N Dimitrova et al. Nature 2008, 1-5 (2008)

Werner Syndrome


Phenotypes

- skin wrinkling
- hair loss and graying
- cataracts
- diabetes (type II)
- osteoporosis
- atherosclerosis, heart disease
- cancer; particularly sarcomas

Caused by loss of the RecQ helicase WRN



Telomere dysfunction contributes to the disease pathology



WS cells exhibit increased:


The shortest telomere, not the average telomere length, is the critical determinant of cell viability and genome stability

(Hemann Cell 2001)



Stochastic telomere loss occurs in normal senescent cells as well

Telomeres Are Hypersensitive to Replication Stress

Fragile Telomere



Telomere loss



Ambrus, NAR 2006

aphidicolin (fragile only)

oncogenic signaling


G-quadruplex ligands

Loss of:

helicases WRN, BLM, RTEL1


shelterin TRF1, POT1, others

Ogg1 and Nth1 glycosylase




Can DNA Damage Cause Telomere Loss or Fragility?

Fragile Telomere






Telomere loss



X = blocking DNA adduct or lesion ??


aphidicolin (fragile only)
oncogenic signaling
G-quadruplex ligands
Loss of:
helicases WRN, BLM, RTEL1
shelterin TRF1, POT1
Ogg1 and Nth1 glycosylase



How Do Telomeres Manage DNA Damage?


DNA Repair Mechanisms

DNA Damage Tolerance



1. Kruk et al. 1995, *PNAS*
2. Rochette & Brash 2010, *PLoS Genetics*

Xeroderma Pigmentosum




- Caused by mutations in genes that code for NER proteins (XP-A to XP-G); or polymerase η (XPV)
- Photosensitivity
- Pigmentation abnormality and atrophic skin
- >1000x increase in skin cancer and 10x increase in internal tumors
- neurological degeneration in 30% of the cases
- Shortened telomeres were observed in sunlight or UVB exposed skin tissue from humans and mice

(Ikeda et al. *Human Pathology*. 2014; Stout & Blasco *Cancer Research*. 2013)

Do DNA lesions cause telomere aberrations ?

Are they repaired at telomeres?




X = UV photoproducts (acute physical exposure)
Cr-DNA adducts (chronic chemical exposure)




Hannah Varsalona-Pope

UV Light and Cr(VI) Exposures Induce Bulky Lesions




5'**TTAGGG**
3'**AATCCC**




DNA Repair & Mutagenesis,
2nd Ed, ASM Press

- removed by NER
- block replicative polymerases
- cause replication dependent breaks
- bypassed by polymerase η
- in yeast for Cr(VI)





Salnikow & Zhitkovich; *Chem. Res. Toxicol.* 2008

Pol η Foci Formation and Localization to Telomeres



Translesion Synthesis is Required for Telomere Preservation After DNA Damage



Summary and Conclusions

- Uncovered new evidence that Pol η protects against Cr(VI)-induced cytotoxicity and replication stress.

- The induction of bulky DNA lesions generates telomere defects consistent with failures in telomere replication – loss and fragility

 - UVC-induced DNA photoproducts

 - Cr(VI)-induced DNA adducts


- Pol η deficient cells show significantly higher levels of UVC- and Cr(VI)-induced telomere defects.

 - Translesion DNA synthesis required for replication of damaged telomeres

Telomeres may be highly dependent on damage bypass mechanisms due to deficiencies in break repair and possible lack of damage removal.


DNA lesions can cause telomere aberrations


Are they repaired at telomeres?




Dhvani Parikh

How Do Telomeres Manage DNA Damage?






Telomere Capture Assay



▲ = pyridimine dimer



11x 100 mm dishes

Telomere < 0.025% of the genome !

20×10^6 cells



100 µg genomic DNA




25 ng telomeric DNA




Dhvani Parikh

Efficiency of Telomere Isolation



- Eluent yields: Mock = 0 ng, Scrambled = 2.6 ng, and Telomere = 10 ng
- Estimated telomere recovery:
bound/ bound + unbound = 33% ± 0.6%



Telomere Southern Blot



Connor Murphy



Purity of Isolated Telomeres

Alu Repeat Blot - ^{32}P (Alu)




Estimated purity: 1.2 ± 0.2 ng non-telomeric DNA/ 10 ng eluent $\rightarrow$ ~90% pure

UVC Exposure Delays Cell Proliferation






- Following UVC exposure cells do not double even after 72 hrs
- We do not expect photoproduct dilution by cell division
- The majority of cells are viable (trypan blue assay)

CPD Repair in Genomic and Telomeric DNA





6-4 PP Repair in Genomic and Telomeric DNA

BJ-hTERT cells




• is telomerase required?

Telomerase is Not Required for Telomere Repair






U2OS cells – telomerase negative





- are photoproducts removed by NER?

NER is Required for 6-4 PP Removal at Telomeres

XP-A skin fibroblasts – NER deficient




UV Photoproduct Inhibits TRF1 Binding



CPD causes a 13-fold inhibition in TRF1 binding


Role for Nucleotide Excision Repair At Telomeres After Damage



Normal fibroblasts
(XPA complemented)


XP20S-derived cells
SV40 transformed

XP-A fibroblasts
(XPA^{-/-})




- UVC irradiation induces a near 2-fold increase in telomere loss and doublets
- Similar results in NER proficient and deficient cells?

XPA Complemented Cells Repair Slowly



α-XPA



α-GAPDH

Elise
Fouquerel

Summary and Future Directions

1. UVC induces CPD and 6-4 PP formation at telomeres
 - frequency is about 2-fold less than the bulk genome
2. CPDs are repaired 1.5-fold faster at telomeres compared to the bulk genome.
Transcription coupled repair?
3. 6-4 PPs persist at telomeres and the bulk genome in XPA deficient cells
 - Indicates that NER is active at telomeres
4. 6-4 PPs removal rates at telomeres are similar to the bulk genome in telomerase positive (BJ-hTERT) and telomerase negative (U2OS) cells
5. UVC generates telomere defects consistent with failures in telomere replication
 - Aberrations are increased in translesion synthesis deficient cells
 - NER deficient cells?

Acknowledgements



Opresko lab

- Sari Joshowitz
- Melinda Sager
- Elise Fouquerel
- Justin Lormand
- Hannah Pope-Varsalona
- Connor Murphy
- Dhvani Parikh

Collaborators

- Hong Wang
- Ben Van Houten
- Rob Sobol
- Simon Watkins (CBI)
- Woody Wright
- Bruce Armitage
- Danith Ly
- Marcel Bruchez

Funding

- R01 ES022944
- R21 AG045545
- R34 GM108187/ PNA Innovations
- David Scaife Foundation CNAIST
- UPCI Skin Cancer SPORE; CDA

UPCI Genome Stability Group
CNAIST (CMU)

Postdoctoral positions available